

Azopt



Azopt reduces the amount of fluid in the eye, which decreases pressure inside the eye. **Azopt** ophthalmic (for the eyes) is used to treat certain types of glaucoma and other causes of high pressure inside the eye.

Active ingredient: [Brinzolamide](#)

Brand(s): Azopt

Manufacturer: Alcon Inc.

Disease(s): Open-angle Glaucoma / Ocular Hypertension

Azopt 1% 5ml

| | Per bottle | Per Pack | Order |
|-----------|------------|-----------------------------------------|-----------------------------|
| 1 bottles | \$ 54.95 | \$ 54.95 | Add to cart |
| 2 bottles | \$ 49.98 | \$ 99.95 \$109.90 | Add to cart |
| 3 bottles | \$ 43.32 | \$ 129.95 \$164.85 | Add to cart |

Brinzolamide drops

What is this medicine?

BRINZOLAMIDE reduces the amount of fluid in the eye, which decreases pressure inside the eye. Brinzolamide ophthalmic (for the eyes) is used to treat certain types of glaucoma and other causes of high pressure inside the eye. Brinzolamide ophthalmic may also be used for other purposes not listed in this medication guide.

What should my health care professional know before I take this medicine?

You should not use this medication if you are allergic to brinzolamide. Before using this medication, tell your doctor if you are allergic to any drugs, or if you have narrow-angle glaucoma. FDA pregnancy category C. This medication may be harmful to an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant during treatment. It is not known whether brinzolamide ophthalmic passes into breast milk or if it could harm a nursing baby. Do not use this medication without telling your doctor if you are breast-feeding a baby.

How should I take this medicine?

Do not use this medication while you are wearing contact lenses. This medication may contain a preservative that can be absorbed by soft contact lenses. Wait at least 15 minutes after using the eye drops before putting your contact lenses in.

Use brinzolamide ophthalmic exactly as it was prescribed for you. Do not use the medication in larger amounts, or use it for longer than recommended by your doctor. Follow the instructions on your prescription label.

Wash your hands before using the eye drops.

To apply the eye drops:

- Tilt your head back slightly and pull down on the lower eyelid to create a small pocket. Hold the dropper above the eye with the dropper tip down. Look up and away from the dropper. Squeeze out a drop and close your eye. Gently press your finger to the inside corner of the eye (near the nose) for about 1 minute to keep the

