

Please click price below and you will be redirected to our pharmacy.

## Combivir



**Generic Combivir** is in a group of medicines called reverse transcriptase inhibitors. It is used for treating HIV infection in combination with other medicines.

Active ingredient: [Lamivudine - Zidovudine](#)

**Brand(s):**Duovir

**Manufacturer:** Cipla Limited

**Disease(s):**Hiv

### Combivir 150mg + 300mg

	Per pill	Per Pack	Order
30 tablet	\$ 2.00	<b>\$ 60.00</b>	<a href="#">Add to cart</a>
60 tablet	\$ 1.83	<b>\$ 110.00</b> <del>\$120.00</del>	<a href="#">Add to cart</a>
90 tablet	\$ 1.67	<b>\$ 150.00</b> <del>\$180.00</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
120 tablet	\$ 1.58	<b>\$ 190.00</b> <del>\$240.00</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
180 tablet	\$ 1.44	<b>\$ 260.00</b> <del>\$360.00</del>	<a href="#">Add to cart</a>

Free AirMail shipping

## Lamivudine and Zidovudine tablet

### What is this medicine?

LAMIVUDINE; ZIDOVUDINE is two antiretroviral drugs in one tablet. It is used with other medicines to treat HIV. This medicine is not a cure for HIV. It will not stop the spread of HIV to others.

### What should my health care professional know before I take this medicine?

They need to know if you have any of these conditions:

- blood or bone marrow problems
- kidney disease
- liver disease
- an unusual or allergic reaction to lamivudine, zidovudine, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I take this medicine?

Take this medicine by mouth with a glass of water. You may take this medicine with or without food. Take your medicine at regular intervals. Do not take your medicine more often than directed. For your anti-HIV therapy to work as well as possible, take each dose exactly as prescribed. Do not skip doses or stop your medicine even if you feel better. Skipping doses may make the HIV virus resistant to this medicine and other medicines. Do not stop taking except on your doctor's advice.

Talk to your pediatrician regarding the use of this medicine in children.

