

Please click price below and you will be redirected to our pharmacy.

## Cialis Oral Jelly



Cialis Oral Jelly is an Erectile Dysfunction drug made in a form of jelly, sorted in a compact sachet. In comparison to other ED products it dissolves much faster that's why starts acting already in 20 minutes, moreover it has a pleasant taste.

Active ingredient: [Tadalafil](#)

**Brand(s):**Tadarise

**Manufacturer:** Sunrise Remedies Pvt. Ltd.

**Disease(s):**Erectile Dysfunction / Impotence

### Cialis Oral Jelly 20mg

	Per jelly	Per Pack	Order
7 jelly	\$ 4.99	<b>\$ 34.95</b>	<a href="#">Add to cart</a>
14 jelly	\$ 4.28	<b>\$ 59.95</b> <del>\$69.90</del>	<a href="#">Add to cart</a>
21 jelly	\$ 3.81	<b>\$ 79.95</b> <del>\$104.85</del>	<a href="#">Add to cart</a>
Viagra Soft 100mg x 4 pills FREE bonus!			
28 jelly	\$ 3.57	<b>\$ 99.95</b> <del>\$139.80</del>	<a href="#">Add to cart</a>
Viagra Soft 100mg x 4 pills FREE bonus!			
35 jelly	\$ 3.43	<b>\$ 119.95</b> <del>\$174.75</del>	<a href="#">Add to cart</a>
Viagra Soft 100mg x 4 pills FREE bonus!			
42 jelly	\$ 3.21	<b>\$ 134.95</b> <del>\$209.70</del>	<a href="#">Add to cart</a>
Viagra Soft 100mg x 4 pills FREE bonus!			
49 jelly	\$ 3.06	<b>\$ 149.95</b> <del>\$244.65</del>	<a href="#">Add to cart</a>
Viagra Soft 100mg x 4 pills FREE bonus!			

## Tadalafil jelly

### What is this medicine?

TADALAFIL is used to treat erection problems in men. Also, it is currently in Phase 3 clinical trials for treating pulmonary arterial hypertension.

### What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- eye or vision problems, including a rare inherited eye disease called retinitis pigmentosa
- heart disease, angina, a history of heart attack, irregular heart beats, or other heart problems
- high or low blood pressure
- kidney or liver disease
- stroke
- an unusual or allergic reaction to tadalafil, other medicines, foods, dyes, or preservatives

### How should I use this medicine?

Take this medicine by mouth with a glass of water. You may take this medicine with or without meals. The dose is usually taken 20 to 60

