

Please click price below and you will be redirected to our pharmacy.

Acofide



Generic Acofide(Acotiamide) is used in the treatment of functional dyspepsia. It helps to relieve the symptoms like bloating after meal, epigastric pain/discomfort and early satiety. Acotiamide increases the release of acetylcholine, a chemical that can increase the motility of the intestine.

Active ingredient: [Acotiamide](#)

Brand(s):Actnew

Manufacturer: Abbott

Disease(s):Functional Dyspepsia / Bloating After Meal / Epigastric Pain / Discomfort

Acofide 100mg

| | Per pill | Per Pack | Order |
|-----------|----------|---|-----------------------------|
| 10 tablet | \$ 5.00 | \$ 49.95 | Add to cart |
| 20 tablet | \$ 4.50 | \$ 89.95 \$99.90 | Add to cart |
| 30 tablet | \$ 4.00 | \$ 119.95 \$149.85 | Add to cart |
| 60 tablet | \$ 3.00 | \$ 179.95 \$299.70 | Add to cart |

Free AirMail shipping

Acotiamide tablets

What is this medicine?

Acotiamide is used in the treatment of functional dyspepsia. It helps to relieve the symptoms like bloating after meal, epigastric pain/discomfort and early satiety. Acotiamide increases the release of acetylcholine, a chemical that can increase the motility of the intestine.

Important information

Before using this medicine, be sure to tell your doctor and pharmacist:

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

How should I take this medicine?

If you miss a dose, skip the missed dose and follow your regular dosing schedule. You should never take two doses at one time.

If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.

Do not stop taking this medicine unless your doctor instructs you to do so.

What side effects may I notice from this medicine?

The most commonly reported adverse reactions include diarrhoea, constipation, nausea, vomiting, rash and hives. If any of these symptoms

