

Please click price below and you will be redirected to our pharmacy.

Noroxin



Generic Norfloxacin is an antibiotic in a group of drugs called fluoroquinolones. It is used to treat bacterial infections of genito-urinary tract.

Active ingredient: [Norfloxacin](#)

Brand(s):Norflox

Manufacturer: Okasa Pharma Pvt. Ltd.

Disease(s):Urinary Tract Infections

Noroxin 400mg

| | Per pill | Per Pack | Order |
|-----------------------|----------|---|-----------------------------|
| 30 tablet | \$ 1.33 | \$ 39.99 | Add to cart |
| 60 tablet | \$ 1.17 | \$ 69.99 \$79.98 | Add to cart |
| 90 tablet | \$ 1.11 | \$ 99.99 \$119.97 | Add to cart |
| 120 tablet | \$ 1.00 | \$ 119.99 \$159.96 | Add to cart |
| 180 tablet | \$ 0.94 | \$ 169.99 \$239.94 | Add to cart |
| Free AirMail shipping | | | |
| 360 tablet | \$ 0.89 | \$ 319.99 \$479.88 | Add to cart |

Free AirMail shipping

Norfloxacin tablet

What is this medicine?

NORFLOXACIN is a quinolone antibiotic. It can kill certain bacteria or stop their growth. This medicine is used to treat infections of the prostate and urinary tract. It is also used to treat some sexually transmitted diseases. It will not work for colds, flu, or other viral infections.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- cerebral disease
- glucose-6- phosphate dehydrogenase related disorder
- history of tendon rupture
- kidney disease
- myasthenia gravis
- seizure disorder
- an unusual or allergic reaction to norfloxacin, other quinolone antibiotics, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

How should I use this medicine?

Take this medicine by mouth with a full glass of water. Take this medicine on an empty stomach, at least 1 hour before or 2 hours after a meal, milk and/or other dairy products. Do not take with food. Take your medicine at regular intervals. Do not take your medicine more often than directed.

Take all of your medicine as directed even if you think your are better. Do not skip doses or stop your medicine early.

