

Please click price below and you will be redirected to our pharmacy.

## Minocin



**Generic Minocin** is a tetracycline antibiotic. It is used to treat many different bacterial infections, such as urinary tract infections, severe acne, gonorrhea, tick fever, chlamydia and others.

Active ingredient: [Minocycline](#)

**Brand(s):**CNN

**Manufacturer:** Ipca Laboratories

**Disease(s):**Infections / Acne / Gonorrhea / Tick Fever / Chlamydia

### Minocin 100mg

	Per pill	Per Pack	Order
30 tablet	\$ 3.63	<b>\$ 108.92</b>	<a href="#">Add to cart</a>
60 tablet	\$ 3.18	<b>\$ 190.71</b> <del>\$217.84</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
90 tablet	\$ 3.15	<b>\$ 283.54</b> <del>\$326.76</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
120 tablet	\$ 2.81	<b>\$ 337.20</b> <del>\$435.68</del>	<a href="#">Add to cart</a>
Free AirMail shipping			

## Minocycline Hydrochloride tablet

### What is this medicine?

MINOCYCLINE is a tetracycline antibiotic. It stops the growth of some bacteria. It is used to treat many kinds of infections, like acne, respiratory, urinary tract, and sexually transmitted infections. It will not work for colds, flu, or other viral infections.

### What should my health care professional know before I take this medicine?

They need to know if you have any of these conditions:

- kidney disease
- liver disease
- an unusual or allergic reaction to minocycline, tetracycline antibiotics, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I take this medicine?

Take this medicine by mouth with a full glass of water. Follow the directions on the prescription label. Take this medicine at least 1 hour before or 2 hours after meals. If it upsets your stomach, you can take it with food. Take your medicine at regular intervals. Do not take your medicine more often than directed. Take all of your medicine as directed even if you think you are better. Do not skip doses or stop your medicine early.

Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

