Xalatan



Generic Xalatan reduces pressure in the eye by increasing the amount of fluid that drains from the eye. Generic Xalatan (for the eyes) is used to treat certain types of glaucoma and other causes of high pressure inside the eye.

Active ingredient: Latanoprost

Brand(s):Latoprost

Manufacturer: Sun Pharmaceutical Industries Ltd.

Disease(s):Glaucoma / Eye Pressure

Xalatan 0.005% 2.5ml

| | Per pill | Per Pack | Order |
|-------------|----------|---------------------------------|-------------|
| 1 eye drops | \$ 39.95 | \$ 39.95 | Add to cart |
| 2 eye drops | \$ 34.98 | \$ 69.95 \$79.90 | Add to cart |
| 3 eye drops | \$ 29.98 | \$ 89.95 \$119.85 | Add to cart |

Latanoprost eye drops

What is this medicine?

LATANOPROST reduces pressure in the eye by increasing the amount of fluid that drains from the eye.

Latanoprost (for the eyes) is used to treat certain types of glaucoma and other causes of high pressure inside the eye.

What should I tell my health care provider before I take this medicine?

To make sure you can safely use Latanoprost, tell your doctor if you have swelling or infection of your eye.

Latanoprost may cause a gradual change in the color of your eyes or eyelids and lashes, as well as increased growth or thickness of your eyelashes. These color changes, usually an increase in brown pigment, occur slowly and you may not notice them for months or years. Color changes may be permanent even after your treatment ends, and may occur only in the eye being treated. This could result in a cosmetic difference in eye or eyelash color from one eye to the other.

FDA pregnancy category C. It is not known whether Latanoprost will harm an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant while using this medication. It is not known whether latanoprost passes into breast milk or if it could harm a nursing baby. Do not use Latanoprost without telling your doctor if you are breast-feeding a baby.

How should I take this medicine?

Do not use Latanoprost while wearing contact lenses. This medication may contain a preservative that can discolor soft contact lenses. Wait at least 15 minutes after using Latanoprost before putting your contact lenses in.

Use Latanoprost exactly as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended. Follow the directions on your prescription label.

Wash your hands before using the Latanoprost eye drops.

To apply the eye drops: