

Please click price below and you will be redirected to our pharmacy.

## Combivent



Combivent is a metered-dose inhaler containing a combination of albuterol and ipratropium. Albuterol and ipratropium are bronchodilators that relax muscles in the airways and increase air flow to the lungs.

Active ingredient: Ipratropium Bromide & Albuterol (salbutamol)

**Brand(s):**Duolin  
**Manufacturer:** Cipla Limited  
**Disease(s):**Chronic Obstructive Pulmonary Disease

	Per item	Per Pack	Order
1 inhaler	\$ 69.95	<b>\$ 69.95</b>	<a href="#">Add to cart</a>
2 inhaler	\$ 59.98	<b>\$ 119.95</b> \$139.90	<a href="#">Add to cart</a>
3 inhaler	\$ 56.65	<b>\$ 169.95</b> \$209.85	<a href="#">Add to cart</a>

Free AirMail shipping

### Albuterol and Ipratorium inhalation

#### What is this medicine?

Albuterol and Ipratorium is used as an inhaled medication to prevent bronchospasm in people with chronic obstructive pulmonary disease (COPD) who are also using other medicines to control their condition. Albuterol and Ipratorium may also be used for purposes not listed in this medication guide.

#### What should I tell my health care provider before I take this medicine?

You should not use Albuterol and Ipratorium if you are allergic to albuterol (Proventil, Ventolin), ipratropium (Atrovent), or atropine (Atreza, Lomotil, Vi-Atro, Sal-Tropine).

To make sure Albuterol and Ipratorium is safe for you, tell your doctor about your other medical conditions, especially:

- heart disease, high blood pressure, coronary artery disease, or heart rhythm disorder;
- a seizure disorder such as epilepsy;
- diabetes;
- overactive thyroid;
- glaucoma;
- enlarged prostate, problems with urination; or
- liver or kidney disease.

FDA pregnancy category C. It is not known whether Albuterol and Ipratorium is harmful to an unborn baby. Tell your doctor if you are pregnant or plan to become pregnant during treatment.

#### How should I use this medicine?

Use Albuterol and Ipratorium exactly as prescribed by your doctor. Do not use in larger or smaller amounts or for longer than recommended. Follow the directions on your prescription label.

