Please click price below and you will be redirected to our pharmacy.

Betadine

Betadine cream is used for treating minor wounds and infections, as well as killing bacteria. **Betadine** cream is an antiseptic combination. It works by killing sensitive bacteria.

Active ingredient: Providone-iodine

Brand(s): Wokadine
Manufacturer: Wockhardt Ltd.
Disease(s): Wound / Infection

Betadine 10% 15g

	Per pill	Per Pack	Order
1 tube	$ 24.95	$ 24.95	Add to cart
2 tube	$ 19.98	$ 39.95 $49.90	Add to cart
3 tube	$ 17.65	$ 52.95 $74.85	Add to cart
4 tube	$ 14.99	$ 59.95 $99.80	Add to cart
5 tube	$ 13.99	$ 69.95 $124.75	Add to cart

Povidone-Iodine topical

What is this medicine?

Povidone-Iodine cream is used for treating minor wounds and infections, as well as killing bacteria. Povidone-Iodine cream is an antiseptic combination. It works by killing sensitive bacteria.

What should I tell my health care providers before I take this medicine?

Some medical conditions may interact with Povidone-Iodine cream. Tell your doctor or pharmacist if you have any medical conditions, especially if any of the following apply to you:

- if you are pregnant, planning to become pregnant, or are breast-feeding
- if you are taking any prescription or nonprescription medicine, herbal preparation, or dietary supplement
- if you have allergies to medicines, foods, or other substances
- if you have serious burns or deep puncture wounds

Some MEDICINES MAY INTERACT with Povidone-Iodine cream. However, no specific interactions are known at this time.

This may not be a complete list of all interactions that may occur. Ask your health care provider if Povidone-Iodine cream may interact with other medicines that you take. Check with your health care provider before you start, stop, or change the dose of any medicine.

How should I use this medicine?

Use Povidone-Iodine cream as directed by your doctor. Check the label on the medicine for exact dosing instructions.

- Clean the affected area. Apply a small amount of Povidone-Iodine cream to the affected area. The area may be left uncovered, or you may cover it with a sterile bandage.
- If you miss a dose of Povidone-Iodine cream, use it as soon as