

Please click price below and you will be redirected to our pharmacy.

## Syphilis test



**Syphilis Test** is a qualitative membrane device based immunoassay for the detection of TP antibodies (IgG and IgM) in whole blood, serum or plasma. In this test procedure, recombinant Syphilis antigen is immobilized in the test line region of the device. After a specimen is added to the specimen well of the device, it reacts with Syphilis antigen coated particles in the test. This mixture migrates chromatographically along the length of the test strip and interacts with the immobilized Syphilis antigen. The double antigen test format can detect both IgG and IgM in specimens. If the specimen contains TP antibodies, a coloured line will appear in the test line region, indicating a positive result. If the specimen does not contain TP antibodies, a coloured line will not appear in this region, indicating a negative result.

Active ingredient: [Syphilis Test](#)

Disease(s): Syphilis

### Syphilis test 1mg

	Per device	Per Pack	Order
1 device	\$ 2.00	<b>\$ 2.00</b>	<a href="#">Add to cart</a>

## Syphilis Rapid Test Device

### What is this product?

**Syphilis Test** is a qualitative membrane device based immunoassay for the detection of TP antibodies (IgG and IgM) in whole blood, serum or plasma. In this test procedure, recombinant Syphilis antigen is immobilized in the test line region of the device. After a specimen is added to the specimen well of the device, it reacts with Syphilis antigen coated particles in the test. This mixture migrates chromatographically along the length of the test strip and interacts with the immobilized Syphilis antigen. The double antigen test format can detect both IgG and IgM in specimens. If the specimen contains TP antibodies, a coloured line will appear in the test line region, indicating a positive result. If the specimen does not contain TP antibodies, a coloured line will not appear in this region, indicating a negative result. To serve as a procedural control, a coloured line will always appear in the control line region, indicating that proper volume of specimen has been added and membrane wicking has occurred.

