

## Avodart



**Generic Avodart** is used for treating symptoms of benign prostatic hyperplasia (BPH) or enlargement of the prostate gland. It also helps to reduce the risk of urinary blockage and the need for prostate surgery.

Active ingredient: [Dutasteride](#)

**Brand(s):**Duprostat / Veltride / Dutanol

**Manufacturer:** Cipla Limited / Intas Pharmaceuticals Ltd. / Knoll Healthcaare Pvt. Ltd.

**Disease(s):**Benign Prostatic Hyperplasia

### Avodart 0.5mg

	Per pill	Per Pack	Order
30 tablet	\$ 1.33	<b>\$ 39.95</b>	<a href="#">Add to cart</a>
60 tablet	\$ 1.25	<b>\$ 74.95</b> <del>\$79.90</del>	<a href="#">Add to cart</a>
90 tablet	\$ 1.11	<b>\$ 99.95</b> <del>\$119.85</del>	<a href="#">Add to cart</a>
120 tablet	\$ 1.00	<b>\$ 119.95</b> <del>\$159.80</del>	<a href="#">Add to cart</a>
180 tablet	\$ 0.92	<b>\$ 164.95</b> <del>\$239.70</del>	<a href="#">Add to cart</a>
Free AirMail shipping			
360 tablet	\$ 0.83	<b>\$ 299.95</b> <del>\$479.40</del>	<a href="#">Add to cart</a>

Free AirMail shipping

## Dutasteride capsule

### What is this medicine?

DUTASTERIDE is used to treat benign prostatic hyperplasia (BPH) in men. This is a condition that causes you to have an enlarged prostate. This medicine helps to control your symptoms, decrease urinary retention, and reduces your risk of needing surgery.

### What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- liver disease
- prostate cancer
- an unusual or allergic reaction to dutasteride, finasteride, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I take this medicine?

Take this medicine by mouth with a glass of water. Do not cut, crush, chew or open this medicine. You can take this medicine with or without food. Take your doses at regular intervals. Do not take your medicine more often than directed. Do not stop taking except on your doctor's advice.

Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdose: If you think you have taken too much of this medicine

