Cialis®



Cialis® is in a class of drugs called Phosphodiesterase Enzyme Inhibitors. It is used to treat erection problems in men. Cialis® is manufactured by Eli Lilly.

Active ingredient: Tadalafil

Brand(s):Cialis®

Manufacturer: Eli Lilly And Company
Disease(s):Erectile Dysfunction / Impotence

Cialis® 20mg

| | Per pill | Per Pack | Order | | | |
|-----------------------|----------|----------------------------------|-------------|--|--|--|
| 8 tablet | \$ 13.74 | \$ 109.95 | Add to cart | | | |
| 12 tablet | \$ 11.66 | \$ 139.95 \$164.93 | Add to cart | | | |
| 20 tablet | \$ 11.00 | \$ 219.95 \$274.88 | Add to cart | | | |
| Free AirMail shipping | | | | | | |
| 32 tablet | \$ 10.31 | \$ 329.95 \$439.80 | Add to cart | | | |
| Free AirMail shipping | | | | | | |
| 60 tablet | \$ 9.67 | \$ 579.95 \$824.62 | Add to cart | | | |
| Free AirMail shipping | | | | | | |

Tadalafil tablet (Cialis®)

What is this medicine?

TADALAFIL is used to treat erection problems in men. It works by dilating blood vessels in the penis, allowing the inflow of blood needed for an erection. Cialis®'s 36-hour effectiveness earned it the nickname, "The Weekend Pill". Also, Cialis® is currently in Phase 3 clinical trials for treating pulmonary arterial hypertension.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- eye or vision problems, including a rare inherited eye disease called retinitis pigmentosa
- heart disease, angina, a history of heart attack, irregular heart beats, or other heart problems
- · high or low blood pressure
- · kidney or liver disease
- stroke
- an unusual or allergic reaction to tadalafil, other medicines, foods, dyes, or preservatives

How should I use this medicine?

Take this medicine by mouth with a glass of water. You may take this medicine with or without meals. The dose is usually taken 30 to 60 minutes before sexual activity. You should not take this dose more than once per day. Do not take your medicine more often than directed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.