

Please click price below and you will be redirected to our pharmacy.

## Fasigyn



Tinidazole is used to treat certain infections caused by bacteria, such as infection of the intestines or vagina. It is also used to treat certain sexually transmitted infections.

Active ingredient: [Tinidazole](#)

**Brand(s):** Tinima / Tinamide

**Manufacturer:** Morepen Laboratories / Ranbaxy Laboratories Ltd.

**Disease(s):** Infection Of The Intestines Or Vagina / Sexually Transmitted Infections

### Fasigyn 500mg

	Per pill	Per Pack	Order
30 tablet	\$ 1.50	<b>\$ 44.95</b>	<a href="#">Add to cart</a>
60 tablet	\$ 1.33	<b>\$ 79.95</b> <del>\$89.90</del>	<a href="#">Add to cart</a>
90 tablet	\$ 1.11	<b>\$ 99.95</b> <del>\$134.85</del>	<a href="#">Add to cart</a>
120 tablet	\$ 1.00	<b>\$ 119.95</b> <del>\$179.80</del>	<a href="#">Add to cart</a>
180 tablet	\$ 0.95	<b>\$ 170.95</b> <del>\$269.70</del>	<a href="#">Add to cart</a>

Free AirMail shipping

## Tinidazole tablets

### What is this medicine?

Tinidazole is an antibiotic that fights bacteria in the body.

Tinidazole is used to treat certain infections caused by bacteria, such as infection of the intestines or vagina. It is also used to treat certain sexually transmitted infections.

Tinidazole may also be used for purposes other than those listed in this medication guide.

### What should I tell my health care provider before I take this medicine?

You should not use this medication if you are allergic to tinidazole or metronidazole (Flagyl), or if you are in the first 3 months of pregnancy.

Tinidazole can pass into breast milk and may harm a nursing baby. **Do not breast-feed while you are taking tinidazole and for at least 3 days after your last dose.**

You may begin nursing again 3 days after your last dose of tinidazole. If you use a breast pump during treatment, throw out any milk you collect while taking tinidazole. Do not feed it to your baby.

To make sure you can safely take tinidazole, tell your doctor if you have any of these other conditions:

- kidney disease (or if you are on dialysis);
- epilepsy or other seizure disorder;
- a blood cell disorder such as anemia or low platelets; or
- a weak immune system.

FDA pregnancy category C. **Do not take tinidazole during the first 3 months of pregnancy.** Tell your doctor if you are pregnant or plan to become pregnant during treatment.

