Boniva



Generic Boniva is used for preventing and treating osteoporosis (weak bones) in women who are past menopause.

Active ingredient: Ibandronae Sodium

Brand(s):Bandrone

Manufacturer: Natco Pharma Limited

Disease(s):Osteoporosis

Boniva 50mg

	Per pill	Per Pack	Order			
10 tablet	\$ 24.99	\$ 249.95	Add to cart			
Free AirMail shipping						
20 tablet	\$ 22.50	\$ 449.95 \$499.90	Add to cart			
Free AirMail shipping						
30 tablet	\$ 20.00	\$ 599.95 \$749.85	Add to cart			
Free AirMail ship	ping					

rice Alliviali Silippilig

Ibandronate Sodium tablet

What is this medicine?

IBANDRONATE slows calcium loss from bones. It is used to treat osteoporosis in women past the age of menopause.

What should I tell my health care provider before I take this medicine?

They need to know if you have any of these conditions:

- · dental disease
- · esophagus, stomach, or intestine problems, like acid reflux or GERD
- · kidney disease
- · low blood calcium
- low vitamin D
- · problems sitting or standing
- · trouble swallowing
- an unusual or allergic reaction to ibandronate, other medicines, foods, dyes, or preservatives
- · pregnant or trying to get pregnant
- · breast-feeding

How should I use this medicine?

You must take this medicine exactly as directed or you will lower the amount of medicine you absorb into your body or you may cause yourself harm. 50mg is a once-a-month supply. Take your dose by mouth first thing in the morning, after you are up for the day. Do not eat or drink anything before you take this medicine. Swallow the tablet with a full glass (6 to 8 ounces) of plain water. Do not take this medicine with any other drink. Do not chew or crush the tablet. After taking this medicine, do not eat breakfast, drink, or take any other medicines or vitamins for at least 1 hour. Stand or sit up for at least 1 hour after taking this medicine; do not lie down. Do not take your medicine more often than directed.

Talk to your pediatrician regarding the use of this medicine in children.