

Please click price below and you will be redirected to our pharmacy.

## Ticlid



Generic **Ticlid** is used for reducing the risk of stroke. It may also be used in combination with aspirin to prevent a blood clot from forming after a stent is placed in the coronary artery.

Active ingredient: **Ticlopidine**

**Brand(s):**Ticlop

**Manufacturer:** Zydus Cadila Healthcare

**Disease(s):**Stroke / Blood Clots

### Ticlid 250mg

|                       | Per pill | Per Pack                                | Order                       |
|-----------------------|----------|---|-----------------------------|
| 30 tablet             | \$ 2.33  | <b>\$ 69.95</b>                         | <a href="#">Add to cart</a> |
| 60 tablet             | \$ 2.17  | <b>\$ 129.95</b><br><del>\$139.90</del> | <a href="#">Add to cart</a> |
| 90 tablet             | \$ 2.11  | <b>\$ 189.95</b><br><del>\$209.85</del> | <a href="#">Add to cart</a> |
| Free AirMail shipping |          |   |                             |
| 120 tablet            | \$ 2.08  | <b>\$ 249.95</b><br><del>\$279.80</del> | <a href="#">Add to cart</a> |
| Free AirMail shipping |          |   |                             |
| 180 tablet            | \$ 2.09  | <b>\$ 376.80</b><br><del>\$419.70</del> | <a href="#">Add to cart</a> |

Free AirMail shipping

## Ticlopidine tablet

### What is this medicine?

TICLOPIDINE helps to prevent blood clots. This medicine is used to prevent strokes in patients who have already had a stroke, or those who are at high risk of having a stroke. This medicine may also be used in combination with aspirin to prevent blood clots in patients who have had a coronary stent placed.

### What should my health care professional know before I take this medicine?

They need to know if you have any of the following conditions:

- anemia
- high cholesterol
- history of bleeding disorder
- kidney disease
- liver disease
- stomach ulcer
- an unusual or allergic reaction to ticlopidine, other medicines, foods, dyes, or preservatives
- pregnant or trying to get pregnant
- breast-feeding

### How should I take this medicine?

Take this medicine by mouth with a glass of water. Take with food or just after eating to reduce stomach upset. Take your doses at regular intervals. Do not take your medicine more often than directed.

