

Depakene



Valproic acid syrup is used to treat seizure disorders, mental/mood conditions (such as manic phase of bipolar disorder), and to prevent migraine headaches. It works by restoring the balance of certain natural substances (neurotransmitters) in the brain.

Active ingredient: [Valproic Acid](#)

Brand(s): Valprol Syrup / Valparin Syrup

Manufacturer: Intas Biopharmaceuticals / Sanofi India Ltd

Disease(s): Nausea / Vomiting / Sleepiness / Dry Mouth

Depakene 200ml

	Per bottle	Per Pack	Order
1 bottles	\$ 14.95	\$ 14.95	Add to cart

Valproic acid syrup

What is valproic acid?

Valproic acid is used to treat various types of seizure disorders. Valproic acid is sometimes used together with other seizure medications.

Valproic acid is also used to treat manic episodes related to bipolar disorder (manic depression), and to prevent migraine headaches.

Valproic acid may also be used for purposes not listed in this medication guide.

Warnings

Valproic acid can cause liver failure that may be fatal, especially in children under age 2 and in people with liver problems caused by certain genetic disorders.

You should not use valproic acid if you have liver disease, a urea cycle disorder, or a genetic disorder such as Alpers' disease or Alpers-Huttenlocher syndrome.

Do not start or stop taking valproic acid during pregnancy without your doctor's advice. This medicine may harm an unborn baby or cause birth defects, but having a seizure during pregnancy could harm both mother and baby.

Do not use valproic acid to prevent migraine headaches if you are pregnant.

Call your doctor at once if the person taking this medicine has signs of liver or pancreas problems, such as: loss of appetite, upper stomach pain (that may spread to your back), ongoing nausea or vomiting, dark urine, swelling in the face, or jaundice (yellowing of the skin or eyes).

Do not stop using valproic acid without your doctor's advice. Stopping suddenly may cause a serious, life-threatening type of seizure.

Before taking this medicine

You should not use valproic acid if you are allergic to it, or if you have:

- liver disease;
- a urea cycle disorder; or
- a genetic mitochondrial (MYE-toe-KON-dree-al) disorder such as Alpers' disease or Alpers-Huttenlocher syndrome, especially in a child younger than 2 years old.

Valproic acid can cause liver failure that may be fatal, especially in children under age 2 and in people with liver problems caused by a genetic mitochondrial disorder.

Tell your doctor if you have ever had:

